

①



Europäisches Patentamt
European Patent Office
Office européen des brevets

⑪ Veröffentlichungsnummer:

⑪ Publication number:

⑪ Numéro de publication:

0 1 9 5 8 1 8

Internationale Anmeldung veröffentlicht durch die
Weltorganisation für geistiges Eigentum unter der Nummer:

WO 86/01940 (art.158 des EPÜ).

International application published by the World
Intellectual Property Organisation under number:

WO 86/01940 (art.158 of the EPC).

Demande internationale publiée par l'Organisation
Mondiale de la Propriété Intellectuelle sous le numéro:

WO 86/01940 (art.158 de la CBE).



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : H01M 6/14, 4/90	A1	(11) International Publication Number: WO 86/ 01940 (43) International Publication Date: 27 March 1986 (27.03.86)
(21) International Application Number: PCT/US85/01787 (22) International Filing Date: 19 September 1985 (19.09.85) (31) Priority Application Number: 652,362 (32) Priority Date: 19 September 1984 (19.09.84) (33) Priority Country: US (71) Applicant: HONEYWELL INC. [US/US]; Honeywell Plaza, Minneapolis, MN 55408 (US). (72) Inventor: DODDAPANENI, Narayan ; 700 Vance Drive, Glenside, PA 19038 (US). (74) Agent: BLINN, Clyde, C.; Honeywell Inc., Honeywell Plaza, Minneapolis, MN 55408 (US).		(81) Designated States: CH (European patent), DE (European patent), FR (European patent), GB (European patent), NL (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HIGH RATE METAL OXYHALIDE CELLS <div style="text-align: center;"> </div> <p>Li/SOCl₂ 3-PLATE LAB CELLS W/ 0.5mm CATHODES, CONSTANT CURRENT DENSITY OF 9mA/cm², 1.5M LiAlCl₄/SOCl₂, 4.5° C.</p>		
(57) Abstract Improved catalysts for the reduction of oxyhalide depolarizers such as SOCl ₂ and SO ₂ Cl ₂ in active-metal non-aqueous cells are provided which are easier to prepare and less expensive than phthalocyanine complexes and which are generally more stable in the cell environment. These catalysts include certain macrocyclic complexes of Schiff's base prepared by the reaction of an aldehyde with a primary diamine. The preferred embodiments include mononuclear metal macrocyclic complexes made from salicylaldehyde and o-phenylenediamine and binuclear metal macrocyclic complexes of 5-methyl isophthalaldehyde and 1,3-diaminopropane.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GA	Gabon	MR	Mauritania
AU	Australia	GB	United Kingdom	MW	Malawi
BB	Barbados	HU	Hungary	NL	Netherlands
BE	Belgium	IT	Italy	NO	Norway
BG	Bulgaria	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali		
FR	France				

HIGH RATE METAL OXYHALIDE CELLS

BACKGROUND OF THE INVENTION

5 The invention herein described was made in the source of or under a contract or subcontract thereunder with the Department of the Army.

Field of the Invention

10 The present invention relates generally to the field of non-aqueous electrochemical cells and, more particularly to a class of catalysts for the reduction of oxyhalide depolarizers such as SOCl_2 and SO_2Cl_2 in such cells.

15

Description of the Prior Art

 Much work has been done in the field of high energy battery systems utilizing highly reactive anode materials such as alkali or alkaline earth metals in
20 combination with a non-aqueous electrolyte. The electrolyte is normally composed of a solute which is commonly a metal salt or a complex metal salt of the anode metal dissolved in a compatible non-aqueous solvent

depolarizer such as SO_2 , SOCl_2 , or SO_2Cl_2 . An inert cathode, usually of carbon black, is also used in such cells.

5 The potential of electrochemical couples, such as those mentioned above as high rate, high-power batteries, has not been fully realized mainly because of excessive cathode polarization particularly at high discharge rates and low operating temperatures. This
10 causes a deterioration of cell performance after a short period of operation. For this reason, much research has been conducted concerning the development of additives such as catalysts to reduce the cathode depolarizer, thereby improving the rate capability of the cathodes.

15 Some improvement in the rate capability of these carbon cathodes has been achieved by doping them with metals such as platinum or metal halides. However, high amounts of the metals or metal halides are required to
20 effectively increase the rate capability of cells such as Li/SOCl_2 cells. Also, the cell performance of these dopants deteriorates with thinner electrodes and at the higher temperatures required for certain applications. More recently, metal-phthalocyanine complexes have been
25 shown to possess significant catalytic activity for the electrochemical reduction of solvent depolarizers such as


thionyl chloride or sulfural chloride, thereby increasing cell capacity. Examples of such are found in U.S. patents 4,252,875 and 4,405,693. The need still exists for the provision of catalyzed cathodes for cells such as
5 Li/SOCl₂ cells which are stable and effective over a wide temperature range and which are sufficiently stable in the cell to be used in both active and reserve configurations.

10

SUMMARY OF THE INVENTION


By means of the present invention, improved catalysts for the reduction of oxyhalide depolarizers such as SOCl₂ and SO₂Cl₂ are provided which are easier to prepare and less expensive than phthalocyanine complexes
15 and which are generally more stable in the cell environment. These catalysts include certain macrocyclic complexes of Schiff's base prepared by the reaction of an aldehyde with a primary diamine. The preferred embodiments include mononuclear metal macrocyclic complexes
20 made from salicylaldehyde and o-phenylenediamine of a structure such as

25



-4-

and binuclear metal macrocyclic complexes of 5-methyl
isophthalaldehyde and 1,3-diaminopropane having a
structure such as



where M or Me represents a metal selected from the group
10 consisting of iron and cobalt and m and n represent
integers.

BRIEF DESCRIPTION OF THE DRAWINGS

In the drawings:

15

FIGURE 1 is a graphical representation of the
discharge characteristics of a Li/SOCl₂ cell employing
prior art catalysts cobalt phthalocyanine (CoPc) and iron
phthalocyanine (FePc) dissolved in the electrolyte
20 compared with the uncatalyzed baseline.

FIGURE 2 is a graphical representation showing
the discharge characteristics of a Li/SOCl₂ cell using
mononuclear CoSB catalyst compared with the uncatalyzed
25 baseline.

FIGURE 3 is a graphical representation of polarization characteristics of a Li/SOCl_2 cell using binuclear metal complexes FeMac and CoMac of the invention;

5

FIGURE 4 is a graphical representation of the discharge characteristics of a cell of FIGURE 2 at 23°C and 63°C with and without CoSB.

10

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Catalyst Preparation

The syntheses of FeMac and CoMac were carried out using the method of Pilkington et al, Australian Journal of Chemistry Vol. 23, 2225 (1970). The syntheses of CoSB and FeSB were carried out using a method described in Jahnke et al, Topics in Current Chemistry, Vol. 1, 133 (1979)..

20

Example 1

A solution of 5-methyl isophthalaldehyde (0.05 mole) dissolved in the minimum volume of boiling methanol was added to a suspension formed by mixing 1,3-diaminopropane (0.05M) with a saturated solution of cobalt chloride in methanol. The mixture was heated under reflux for 1 hour. Methanol was removed by boiling at


25

atmospheric pressure. The product was poured into ten times its volume of tetrahydrofuran. The product was filtered, washed with water and dried.

5 Example 2

The process of Example 1 was repeated except that iron chloride was substituted for cobalt chloride of Example 1.

10 The process of Examples 1 and 2 produced binuclear metal macrocyclic complexes CoMac and FeMac, respectively, of the following structure:



where M represents iron or cobalt and m and n are integers.

20

Example 3


A solution of salicylaldehyde (0.05M), o-phenylenediamine (0.05M) and cobalt acetate (0.025M) was refluxed in ethanol for about 30 minutes. The resulting product was washed with cold methanol and dried at 120°C.

25

Example 4

Example 3 was repeated substituting iron acetate for the cobalt acetate.

The structure of the complexes formed in Examples 3 and 4 known as CoSB and FeSB, respectively, was as follows:



Performance of Catalyst in Cells

Tables 1 and 2 show a comparison of discharge characteristics of cells using a prior art catalyst, the catalyst of Example 3, and no catalyst.

TABLE 1. DISCHARGE CHARACTERISTICS OF Li/SOCl₂ CELLS WITH VARIOUS CATALYSTS WITH A CONSTANT CURRENT DENSITY OF 30 MA/cm² AT 23°C

Catalyst	Cell Voltage (V)	Discharge Life (Minutes)
None	2.92	34
Iron Phthalocyanine (FePc)	3.30	52
Mononuclear Couples with Schiff's Base (CoSB)	3.30	70

- ELECTROLYTE = (0.5 M LiAlCl₄ + 1.0 M AlCl₃)/SOCl₂
- CATHODE THICKNESS = 1.0 millimeter

TABLE 2. DISCHARGE CHARACTERISTICS OF Li/SOCl₂ CELLS WITH AND WITHOUT COSB COMPLEX AT A CONSTANT CURRENT LOAD OF 10 MA/cm² AT DIFFERENT OPERATING TEMPERATURES

Catalyst	-32°C		-18°C		23°C		63°C	
	Voltage at 50% DOD (V)	DOD (min)	Voltage at 50% DOD (V)	DOD (min)	Voltage at 50% DOD (V)	DOD (min)	Voltage at 50% DOD (V)	DOD (min)
None	2.70	55	2.80	80	3.12	105	3.36	164
CosB	2.72	60	2.94	90	3.39	182	3.44	193

FIGURE 1 shows the discharge characteristics of prior art cells using cobalt phthalocyanine (CoPc) and iron phthalocyanine (FePc) catalysts dissolved in the electrolyte. It should be noted that the FePc exhibits a somewhat higher catalytic activity. FIGURES 2-4 show data derived using the catalysts of the present invention.

In FIGURES 2-4, the discharge characteristics of Li/SOCl₂ cells catalyzed by various metal macrocyclic complexes are compared to an uncatalyzed cell. All catalyzed cells showed substantial improvement in both cell voltage and specific cathode capacity over the uncatalyzed cell. These complexes have shown as good or better catalytic activity for the reduction of SOCl₂ than the iron phthalocyanine catalyst. In the case of metal phthalocyanine catalysts, FePc showed the best

catalytic activity, whereas, cobalt tetraphenylporphyrin (CoTPP) and cobalt Schiff's base complexes (CoSB) possess superior catalytic activity for SOCl_2 reduction than the iron complexes of these macrocyclic rings. Electronic charge density on the central metal ion, extent of orbital overlapping backbonding and the nature of the specific interaction of the complex with the carbon surface, are some of the factors that can influence the catalytic activity of these complexes for the reduction of SOCl_2 . In the case of the binuclear macrocyclic complexes the iron couples appear to be superior.

The operating temperature also appears to influence the catalytic activity as shown in Table 2. In general, the cell performance improved with catalyzed cells irrespective of operating temperatures and current densities. At low operation temperatures, however, the catalytic activity of cobalt Schiff's base complex is rather reduced but the significance of this behavior is not yet understood. However, one of the factors could be the specific interaction of the CoSB complex with carbon cathode surface. A better way of impregnating the complex onto carbon should show better results since the performance of cells with this catalyst is drastically improved at high operating temperatures. The catalyst stability is projected to be responsible for the superior performance at high operating temperatures.

CLAIMS

The embodiments of the invention in which an exclusive property or right is claimed are defined as follows:

- 5 1. In a non-aqueous electrochemical cell system having an active metal anode, a cathode and a non-aqueous electrolyte having an oxyhalide depolarizer, the improvement comprising:
a catalyst for the reduction of said oxyhalide depolar-
10 izer comprising a metal complex of Schiff's base, said Schiff's base being selected from the group consisting of macrocyclic complexes of 5-methyl isophthalaldehyde and a diamino alkane and complexes of salicylaldehyde and o-phenylenediamine, and
15 wherein the metal in said metal complex is selected from the group consisting of iron and cobalt.
2. The cell of claim 1 wherein said metal is cobalt.
- 20 3. The cell of claim 1 wherein said metal is iron.
- 25 4. The cell of claim 1 wherein said Schiff's base is one made from salisaldehyde and o-phenylene-diamine and said metal complex is mononuclear.

5. The cell of claim 3 wherein said metal is cobalt.

5 6. The cell of claim 3 wherein said metal is iron.

10 7. The cell of claim 1 wherein said Schiff's base is one made from 5-methyl isophthalaldehyde and 1,3-diaminopropane and wherein said metal complex is binuclear.

8. The cell of claim 5 wherein said metal is cobalt.

15 9. The cell of claim 5 wherein said metal is iron.

20 10. In a non-aqueous electrochemical cell having a lithium anode, a carbon cathode and a non-aqueous electrolyte comprising LiAlCl_4 salt in SOCl_2 depolarizer solvent the improvement comprising:
a catalyst for the reduction of said oxyhalide depolarizer comprising a metal complex of Schiff's base, said shift base being selected from the group
25 consisting of macrocyclic complexes of 5-methyl

(Claim 10 continued)

isophthalaldehyde and a diamino alkane and complexes
of salicylaldehyde and o-phenylenediamine, and
wherein the metal in said metal complex is selected
5 from the group consisting of iron and cobalt.

11. The cell of claim 10 wherein said metal is
cobalt.

10 12. The cell of claim 10 wherein said metal is
iron.






Fig. 1
(PRIOR ART)

Li/SOCl₂ CELLS W/0.5mm CATHODES, CONSTANT
CURRENT DENSITY OF 10 mA/cm².
1.5 M LiAlCl₄/SOCl₂, 23° C.






Fig. 2

Li/SOCl₂ 3-PLATE LAB CELLS W/0.5mm CATHODES,
CONSTANT CURRENT DENSITY OF 9 mA/cm².
1.5 M LiAlCl₄/SOCl₂, 4.5° C.



INTERNATIONAL SEARCH REPORT

International Application No PCT/US 85/01787

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC
 IPC⁴: H 01 M 6/14; H 01 M 4/90

II. FIELDS SEARCHED

Classification System	Minimum Documentation Searched ⁷
IPC ⁴	H 01 M

Documentation Searched other than Minimum Documentation
 to the extent that such Documents are included in the Fields Searched *

III. DOCUMENTS CONSIDERED TO BE RELEVANT*

Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	US, A, 4469763 (F. WALSH et al.) 4 September 1984, see claim 1; column 4, lines 20-23 --	1
A	DE, A, 2052955 (BATTELLE INSTITUT) 4 May 1972	
A	Extended Abstracts, volume 83, no. 2, 9-14 October 1983, Washington, D.C., (US) F. Walsh et al.: "An improved Li/Soll ₂ battery", abstract no. 33, see pages 54,55 --	
A	GB, A, 2108312 (HONEYWELL INC.) 11 May 1983, see claim 1 -----	1

* Special categories of cited documents: ¹⁰

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Δ" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

19th December 1985

Date of Mailing of this International Search Report

30 JAN. 1986

International Searching Authority

EUROPEAN PATENT OFFICE

Signature of Authorized Officer

G.L.M. Bruydenberg

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON

INTERNATIONAL APPLICATION NO. PCT/US 85/01787 (SA 10888)

This Annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 21/01/86

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4469763	04/09/84	EP-A- 0141209	15/05/85
DE-A- 2052955	04/05/72	None	
GB-A- 2108312	11/05/83	None	

For more details about this annex :
see Official Journal of the European Patent Office, No. 12/82